arXiv:1402.3543v2 [cs.CC] 10 Aug 2015

Inequalities and tail bounds for elementary symmetric polynomials with applications

Parikshit Gopalan* Amir Yehudayoff†

Abstract

We study the extent of independence needed to approximate the product of bounded random variables in expectation, a natural question that has applications in pseudorandomness and min-wise independent hashing.

For random variables whose absolute value is bounded by 1, we give an error bound of the form $\sigma^{\Omega(k)}$ where k is the amount of independence and σ^2 is the total variance of the sum. Previously known bounds only applied in more restricted settings, and were quanitively weaker. We use this to give a simpler and more modular analysis of a construction of min-wise independent hash functions and pseudorandom generators for combinatorial rectangles due to Gopalan *et al.*, which also slightly improves their seed-length.

Our proof relies on a new analytic inequality for the elementary symmetric polynomials $S_k(x)$ for $x \in \mathbb{R}^n$ which we believe to be of independent interest. We show that if $|S_k(x)|, |S_{k+1}(x)|$ are small relative to $|S_{k-1}(x)|$ for some $k > 0$ then $|S_\ell(x)|$ is also small for all $\ell > k$. From these, we derive tail bounds for the elementary symmetric polynomials when the inputs are only k-wise independent.

*Microsoft Research. parik@microsoft.com.

†Department of Mathematics, Technion–IIT. amir.yehudayoff@gmail.com. Horev fellow – supported by the Taub foundation. Research supported by ISF and BSF

1 Introduction

The power of independence in probability and randomized algorithms stems from the fact that it lets us control expectations of products of random variables. If $X_1, \ldots, X_n$ are independent random variables, then $\mathbb{E}[\prod_{i=1}^n X_i] = \prod_{i=1}^n \mu_i$ where μ_i are their respective means. However, there are numerous settings in computer science, where true independence either does not hold, or is too expensive (in terms of memory or randomness). Motivated by this, we explore settings when approximate versions of the product rule for expectations hold even with limited independence. Concretely, let $X_1, \ldots, X_n$ be random variables lying in the range $[-1, 1]$, with mean μ_i and variance σ_i^2 repectively. We are interested in the smallest $k = k(\delta)$ such that whenever the X_is are drawn from a k-wise independent distribution $\mathcal{D}$, it holds that

$$\left| \mathbb{E}_{\mathcal{D}}[\prod_{i=1}^n X_i] - \prod_{i=1}^n \mu_i \right| \leq \delta. \tag{1}$$

As stated, we cannot hope to make do even with $k = n - 1$. Consider the case where each X_i is a random $\{\pm 1\}$ bit. If $X_n = \prod_{i \leq n-1} X_i$, the resulting distribution is $(n-1)$-wise independent, but $\mathbb{E}[\prod_i X_i] = 1$, whereas it is 0 with true independence. So, clearly, we need some additional assumptions about the random variables.

The main message of this paper is that small total variance is sufficient to ensure that the product rule holds approximately even under k-wise independence.

Theorem 1. *Let $X_1, \ldots, X_n$ be random variables each distributed in the range $[-1, 1]$, with mean μ_i and variance σ_i^2 repectively. Let $\sigma^2 = \sum_i \sigma_i^2$. There exist constants $c_1 > 1$ and $1 > c_2 > 0$ such that under any k-wise independent distribution $\mathcal{D}$,*

$$\left| \mathbb{E}_{\mathcal{D}}[\prod_{i=1}^n X_i] - \prod_{i=1}^n \mu_i \right| \leq (c_1 \sigma)^{c_2 k}. \tag{2}$$

Specifically, if $\sigma < 1/(2c_1)$ then $k = O(\log(1/\delta)/\log(1/\sigma))$-wise independence suffices for Equation (1).

An important restriction that naturally arises is positivity, where each X_i lies in the interval $[0, 1]$. This setting of parameters (positive variables, small total variance) is important for the applications considered in this paper: pseudorandom generators for combinatorial rectangles [EGL+98, LLSZ97] and min-wise independent permutations [BCFM00]. The former is an important problem in the theory of unconditional pseudorandomness which has been studied intensively [EGL+98, LLSZ97, SSZZ99, ASWZ96, Lu02, GMR+12]. Min-wise independent hashing was introduced by Broder et al. [BCFM00] motivated by similarity estimation, and further studied by [Ind99, BCM98, SSZZ99]. [SSZZ99] showed that PRGs for rectangles give min-wise independent hash functions.

The results of [EGL+98, Ind99] tell us that under k-wise independence, positivity and boundedness, the LHS of Equation (1) is bounded by $\exp(-\Omega(k))$, hence $k = O(\log(1/\delta))$ suffices for error δ. In contrast, we have seen that such a bound cannot hold in the $[-1, 1]$ case. However, once the variance is smaller than some constant, our bound beats this bound even in the $[0, 1]$ setting. Concretely, when $\sigma^2 < n^{-\varepsilon}$ for some $\varepsilon > 0$, our result says that $O(1)$-wise independence suffices for inverse polynomial error in Equation (1), as opposed to $O(\log(n))$-wise independence. This improvement is crucial in analyzing PRGs and hash functions in

the polynomially small error regime. A recent result of [GMR$^+$12] achieves near-logarithmic seed-length for both these problems, even in the regime of inverse polynomial error. Their construction is simple, but its analysis is not. Using our results, we give a modular analysis of the pseudorandom generator construction for rectangles of [GMR$^+$12], using the viewpoint of hash functions.

Our analysis is simpler and perhaps more intuitive. It also improves the seed-length of the construction, getting the dependence on the dimension n down to $O(\log\log(n))$ as opposed to $O(\log(n))$, which (nearly) matches a lower bound due to [LLSZ97]. Given the basic nature of the question, we feel our results might find other applications. Very recently, [GKM15] constructed the first pseudorandom generators with near-logrithmic seed-length for several classes of functions including halfspaces, modular tests and combinatorial shapes. The key technical ingredient of their work is a generalization of Theorem 1 to the setting where each X_i takes values in the unit complex disc.

The main technical ingredient in our work is a new analytic inequality about symmetric polynomials in real variables which we believe is independently interesting. The k'th symmetric polynomial in $a = (a_1, a_2, \ldots, a_n)$ is defined as

$$S_k(a) = \sum_{T\subseteq[n]:|T|=k} \prod_{i\in T} a_i \tag{3}$$

(we let $S_0(a) = 1$). We show that for any real vector a, if $|S_k(a)|, |S_{k+1}(a)|$ are small relative to $|S_{k-1}(a)|$ for some $k > 0$, then $|S_\ell(a)|$ is also small for all $\ell > k$. This strengthens and generalzies a result of [GMR$^+$12] for the case $k = 1$.

We give an overview of the new inequality, its use in the derivation of bounds under limited independence, and finally the application of these bounds to the construction of pseudorandom generators and hash functions.

1.1 An inequality for elementary symmetric polynomials

The elementary polynomials appear as coefficients of a univariate polynomial with real roots, since $\prod_{i\in[n]}(\xi + a_i) = \sum_{k=0}^{n} \xi^k S_{n-k}(a)$. Symmetric polynomials have been well studied in mathematics, dating back to classical results of Newton and Maclaurin (see [Ste04] for a survey). This work focuses on their growth rates. Specifically, we study how local information on $S_k(a)$ for two consecutive values of k implies global information for all larger values of k.

It is easy to see that symmetric polynomials over the real numbers have the following property:

Fact A. *Over the real numbers, if $S_1(b) = S_2(b) = 0$ then $b = 0$.*

This is equivalent to saying that if $p(\xi)$ is a real univariate polynomial of degree n with n nonzero roots and $p'(0) = p''(0) = 0$ then $p \equiv 0$. This does not hold over all fields, for example, the polynomial $p(\xi) = \xi^3 + 1$ has three nonzero complex roots and $p'(0) = p''(0) = 0$.

A robust version of Fact A was recently proved in [GMR$^+$12]: For every $a \in \mathbb{R}^n$ and $k \in [n]$,

$$|S_k(a)| \le \left(S_1^2(a) + 2|S_2(a)|\right)^{k/2}. \tag{4}$$

That is, if $S_1(a), S_2(a)$ are small in absolute value, then so is everything that follows. We provide an essentially optimal bound.

Theorem 2. *For every $a \in \mathbb{R}^n$ and $k \in [n]$,*

$$|S_k(a)| \leq \left(\frac{6e(S_1^2(a) + |S_2(a)|)^{1/2}}{k^{1/2}} \right)^k .$$

The parameters promised by Theorem 2 are tight up to an exponential in k which is often too small to matter (we do not attempt to optimise the constants). For example, if $a_i = (-1)^i$ for all $i \in [n]$ then $|S_1(a)| \leq 1$ and $|S_2(a)| \leq n + 1$ but $S_k(a)$ is roughly $(n/k)^{k/2}$.

A more general statement than Fact A actually holds (see Appendix A for a proof).

Fact B. *Over the reals, if $S_k(a) = S_{k+1}(a) = 0$ for $k > 0$ then $S_\ell(a) = 0$ for all $\ell \geq k$.*

We prove a robust version of this fact as well: A twice-in-a-row bound on the increase of the symmetric functions implies a bound on what follows.

Theorem 3. *For every $a \in \mathbb{R}^n$, if $S_k(a) \neq 0$ and*

$$\left| \binom{k+1}{k} \frac{S_{k+1}(a)}{S_k(a)} \right| \leq C \quad \text{and} \quad \left| \binom{k+2}{k} \frac{S_{k+2}(a)}{S_k(a)} \right| \leq C^2$$

then for every $1 \leq h \leq n - k$,

$$\left| \binom{k+h}{k} \frac{S_{k+h}(a)}{S_k(a)} \right| \leq \left(\frac{6eC}{h^{1/2}} \right)^h .$$

Theorem 3 is proved by reduction to Theorem 2. The proof of Theorem 2 is analytic and uses the method of Lagrange multipliers, and is different from that of [GMR+12] which relied on the Newton-Girrard identities. The argument is quite general, and similar bounds may be obtained for functions that are recursively defined. The proof can be found in Section 2.

Stronger bounds are known when the inputs are nonnegative. When $a_i \geq 0$ for all $i \in [n]$, the classical Maclaurin inequalities [Ste04] imply that $S_k(a) \leq (e/k)^k (S_1(a))^k$. In contrast, when we do not assume non-negativity, one cannot hope for such bounds to hold under the assumption that $|S_1(a)|$ or any single $|S_k(a)|$ is small (cf. the alternating signs example above).

1.2 Expectations of products under limited independence

We return to the question alluded to earlier about how much independence is required for the approximate product rule of expectation. This question arises in the context of min-wise hashing [Ind99], PRGs for combinatorial rectangles [EGL+98, GMR+12], read-once DNFs [GMR+12] and more.

One could derive bounds of similar shape to ours using the work of [GMR+12], but with much stronger assumptions on the variables. More precisely, one would require $\mathbb{E}[X_i^{2k}] \leq (2k)^{2k} \sigma_i^{2k}$ for all $i \in [n]$, and get an error bound of roughly $k^{O(k)} (\sum_i \sigma_i^2)^{\Omega(k)}$. These stronger assumptions limit the settings where their bound can be applied (biased variables typically do not have good moment bounds), and ensuring these conditions hold led to tedious case analysis in analyzing their PRG construction.

We briefly outline our approach. We start from the results of [EGL+98, Ind99] who give an error bound of $\exp(-k)$. To prove this, they consider random variables $Y_i = 1 - X_i$, so that

$$\prod_{i=1}^{n} X_i = \prod_{i=1}^{n}(1 - Y_i) = \sum_{j=0}^{n}(-1)^j S_j(Y_1, \ldots, Y_n). \tag{5}$$

By inclusion-exclusion/Bonferroni inequalities, the series on the right gives alternating upper and lower bounds, and the error incurred by truncating to k terms is bounded by $S_k(Y)$. So we can bound the expected error by $\mathbb{E}[S_k(Y)]$ for which k-wise independence suffices.

Our approach replaces inclusion-exclusion by a Taylor-series style expansion about the mean, as in [GMR+12]. Let us assume $\mu_i \neq 0$ and let $X_i = \mu_i(1 + Z_i)$. Thus,

$$\prod_{i=1}^{n} X_i = \prod_{i=1}^{n} \mu_i(1 + Z_i) = \prod_{i=1}^{n} \mu_i \left(\sum_{j=0}^{n} S_j(Z) \right). \tag{6}$$

If this series were alternating, then we would only need to bound $\mathbb{E}[|S_k(Z)|]$, which is easy. However, this need not be true since Z may have negative entries (even if we start with X_is all positive). So, to argue that the first k terms are a good approximation, we need to bound the tail $\sum_{\ell \geq k} S_\ell(Z)$. At first, this seems problematic, since this involves high degree polynomials, and it seems hard to get their expectations right assuming just k-wise independence[1]. Even though we cannot bound $\mathbb{E}[S_\ell(Z)]$ under k-wise independence once $\ell \gg k$, we use our new inequalities for symmetric polynomials to get strong tail bounds on them. This lets us show that truncating Equation (6) after k terms gives error roughly $O(\sigma^{ck})$, and thus $k = O(\log(1/\delta)/\log(1/\sigma)$ suffices for error δ. We next describe these tail bounds in detail.

We assume the following setup: $Z = (Z_1, \ldots, Z_n)$ is a vector of real valued random variables where Z_i has mean 0 and variance σ_i^2, and $\sigma^2 = \sum_i \sigma_i^2 < 1$. Let $\mathcal{U}$ denote the distribution where the coordinates of Z are independent. One can show that $\mathbb{E}_{Z \in \mathcal{U}}[|S_\ell(Z)|] \leq \sigma^\ell/\sqrt{\ell!}$ and hence by Markov's inequality (see Corollary 10) when $t > 1$ and $t\sigma \leq 1/2$,

$$\Pr_{Z \in \mathcal{U}}\left[\sum_{\ell=k}^{n} |S_\ell(Z)| \geq 2(t\sigma)^k \right] \leq 2t^{-2k}. \tag{7}$$

Although k-wise independence does not suffice to bound $\mathbb{E}[S_\ell(Z)]$ for $\ell \gg k$, we use Theorem 3 to show that a similar tail bound holds under limited independence.

Theorem 4. *Let $\mathcal{D}$ denote a distribution over $Z = (Z_1, \ldots, Z_n)$ as above where the Z_is are $(2k+2)$-wise independent. For $t > 0$ and[2] $16et\sigma \leq 1$,*

$$\Pr_{X \in \mathcal{D}}\left[\sum_{\ell=k}^{n} |S_\ell(Z)| \geq 2(6et\sigma)^k \right] \leq 2t^{-2k}. \tag{8}$$

Typically proofs of tail bounds under limited independence proceed by bounding the expectation of some suitable low-degree polynomial. The proof of Theorem 4 does not follow this route. In Section 3.1, we give an example of Z_is and a $(2k+2)$-wise independent distribution on where $\mathbb{E}[|S_\ell(Z)|]$ for $\ell \in \{2k+3, \ldots, n-2k-3\}$ is much larger than under the uniform distribution. The same example also shows that our tail bounds are close to tight.

[1] We formally show this in Section 3.1.

[2] A weaker but more technical assumption on t, σ, k suffices, see Equation (24).

1.3 Applications to pseudorandom generators and hash functions

A hash function is a map $h : [n] \to [m]$. Let $\mathcal{U}$ denote the family of all hash functions $h : [n] \to [m]$. Let $\mathcal{H} \subseteq \mathcal{U}$ be a family of hash functions. For $S \subseteq [n]$, let $\min h(S) = \min_{x \in S} h(x)$. The notion of min-wise independent hashing was introduced by Broder et al. [BCFM00] motivated by similarity estimation, and independently by Mulmuley [Mul96] motivated by computational geometry. The following generalization was introduced by Broder et al. [BCM98]:

Definition 5. We say that $\mathcal{H} : [n] \to [m]$ is approximately ℓ-minima-wise independent with error ε if for every $S \subseteq [n]$ and for every sequence $T = (t_1, \ldots, t_\ell)$ of ℓ distinct elements of S,

$$\left| \Pr_{h \in \mathcal{H}}[h(t_1) < \cdots < h(t_\ell) < \min h(S \setminus T)] - \Pr_{h \in \mathcal{U}}[h(t_1) < \cdots < h(t_\ell) < \min h(S \setminus T)] \right| \leq \varepsilon.$$

Combinatorial rectangles are a well-studied class of tests in pseudorandomness [EGL$^+$98, LLSZ97, SSZZ99, ASWZ96, Lu02, GMR$^+$12]. In addition to being a natural class of statistical tests, constructing generators for them with optimal seeds (up to constant factors) will improve on Nisan's generator for logspace [ASWZ96], a long-standing open problem in derandomization.

Definition 6. A combinatorial rectangle is a function $f : [m]^n \to \{0, 1\}$ which is specified by n co-ordinate functions $f_i : [m] \to \{0, 1\}$ as $f(x_1, \ldots, x_n) = \prod_{i \in m} f_i(x_i)$. A map $\mathcal{G} : \{0, 1\}^r \to [m]^n$ is a PRG for combinatorial rectangles with error ε if for every combinatorial rectangle $f : [m]^n \to \{0, 1\}$,

$$\left| \mathbb{E}_{x \in \{0,1\}^r}[f(\mathcal{G}(x))] - \mathbb{E}_{x \in [m]^n}[f(x)] \right| \leq \varepsilon.$$

A generator $\mathcal{G} : \{0, 1\}^r \to [m]^n$ can naturally be thought of as a collection of 2^r hash functions, one for each seed. For $y \in \{0, 1\}^r$, let $\mathcal{G}(y) = (x_1, \ldots, x_n)$. The corresponding hash function is given by $g_y(i) = x_i$. The corresponding hash functions have the property that the probability that they fool all test functions given by combinatorial rectangles. Saks et al. [SSZZ99] showed that this suffices for ℓ-minima-wise independence. They state their result for $\ell = 1$, but their proof extends to all ℓ (see appendix A). Constructions of PRGs for rectangles and min-wise hash functions that achieve seed-length $O(\log(mn) \log(1/\varepsilon))$ were given by [EGL$^+$98] and [Ind99] respectively using limited independence. The first construction $\mathcal{G}_{\mathcal{MR}}$ to achieve seed-length $\tilde{O}(\log(mn/\varepsilon))$ was given recently by [GMR$^+$12]. We use our results to give an anlysis of their generator which we believe is simpler and more intutitive, which also improves the seed-length, to (nearly) match the lower bound from [LLSZ97].

We take the view of $\mathcal{G}_{\mathcal{MR}}$ as a collection of hash functions $g : [n] \to [m]$, based on iterative applications of an *alphabet squaring* step. We describe the generator formally in Section 5. We start by observing that fooling rectangles is easy when m is small; $O(\log(1/\delta))$-wise independnce suffices, and this requires $O(\log(1/\delta) \log(m)) = O(\log(1/\delta))$ random bits for $m = O(1)$.

The key insight in [GMR$^+$12] is that gradually increasing the alphabet is also easy (in that it requires only logarithmic randomness). Assume that we have a hash function $g_0 : [n] \to [m]$ and from it, we define $g_1 : [n] \to [m^2]$. To do this, we pick a function $g_1' : [n] \times [m] \to [m^2]$ and set $g_1(i) = g_1'(i, g_0(i))$. The key observation is that it suffices to pick g_1' using only $O(\log(1/\delta)/\log(m))$-wise independence (rather than the $O(\log(1/\delta))$-wise independence needed for one shot).

To see why this is so, fix subsets $S_i \subset [m^2]$ for each co-ordinate and pretend that g_0 is truly random. One can show that $\Pr_{g_0}[g_1(i) \in S_i]$ is a random variable over the choice of g_1' with

variance $1/\mathrm{poly}(m)$. Since we are interested in $\prod_i \Pr_{g_0}[g_1(i) \in S_i]$, which is the product of n small variance random variables, Theorem 1 says it suffices to use limited independence[3].

Theorem 7. *Let $\mathcal{G}_{\mathcal{MR}}$ be the family of hash functions from $[n]$ to $[m]$ defined in Section 5.1 with error parameter $\delta > 0$. The seed length is at most $O((\log\log(n)+\log(m/\delta))\log\log(m/\delta))$. Then, for every $S_1, \ldots, S_n \subseteq [m]$,*

$$\left| \Pr_{g \in \mathcal{G}_{\mathcal{MR}}}[\forall\ i \in [n]\ g(i) \in S_i] - \Pr_{h \in \mathcal{U}}[\forall\ i \in [n]\ h(i) \in S_i] \right| \leq \delta.$$

This improves the [GMR+12] bound in the dependence on n and δ (their bound was $O(\log(mn/\delta)\log\log(m)+\log(1/\delta)\log\log(1/\delta)\log\log\log(1/\delta))$). In particular, the dependence on n reduces from $\log(n)$ to $\log\log(n)$[4]. [LLSZ97] showed a lower bound of $\Omega(\log(m) + \log(1/\varepsilon) + \log\log(n))$ even for hitting sets, so our bound is tight upto the $\log\log(m/\delta)$ factor. While [LLSZ97] constructed hitting-set generators for rectangles with near-optimal seedlength, we are unaware of previous constructions of pseudorandom generators for rectangles where the dependence of the seedlength on n is $o(\log(n))$.

Combining this with Theorem 19, we get the following corollary.

Corollary 8. *For every ℓ, there is a family of approximately ℓ-minima-wise independent hash functions with error ε and seed length at most $O((\log\log(n) + \log(m^\ell/\varepsilon))(\log\log(m^\ell/\varepsilon)))$.*

1.4 Subsequent work

Very recently, Gopalan, Kane and Meka [GKM15] constructed the first pseudorandom generators with seed-length $O((\log(n/\delta)\log\log(n/\delta)^2)$ for several classes of functions including halfspaces, modular tests and combinatorial shapes. The key technical ingredient of their work is a generalization of Theorem 1 to the setting where the X_is are complex valued random variables lying in the unit disc. Their proof however is very different from ours, and in particular it does not imply the inequalities and tail bounds for symmetric polynomials that are proved here.

Organization: We present the proofs of our inequalities for symmetric polynomials in Section 2 and tail bounds for symmetric polynomials in Section 3. We use these bounds to prove Theorem 1 on products of low-variance variables in Section 4 and to analyze the [GMR+12] generator in Section 5.

2 Inequalities for symmetric polynomials

Proof of Theorem 2. It will be convenient to use

$$E_2(a) = \sum_{i \in [n]} a_i^2.$$

[3]To optimize the seed-length, we actually use almost k-wise independence rather than exact k-wise independence. So the analysis does not use Theorem 1 as a black-box, but rather it directly uses Theorem 4.

[4]The reason $\log\log(n)$ seedlength is possible is because every rectangle can be ε-approximated by one that depends only on $O(m\log(1/\varepsilon))$ co-ordinates. Hence the number of functions to fool grows polynomially in n, rather than exponentially.

By Newton's identity, $E_2 = S_1^2 - 2S_2$ so for all $a \in \mathbb{R}^n$,

$$S_1^2(a) + E_2(a) \leq 2(S_1^2(a) + |S_2(a)|).$$

It therefore suffices to prove that for all $a \in \mathbb{R}^n$ and $k \in [n]$,

$$S_k^2(a) \leq \frac{(16e^2(S_1^2(a) + E_2(a)))^k}{k^k}.$$

We prove this by induction. For $k \in \{1, 2\}$, it indeed holds. Let $k > 2$. Our goal will be upper bounding the maximum of the projectively defined[5] function

$$\phi_k(a) = \frac{S_k^2(a)}{(S_1^2(a) + E_2(a))^k}$$

under the constraint that $S_1(a)$ is fixed. Since ϕ_k is projectively defined, its supremum is attained in the (compact) unit sphere, and is therefore a maximum. Choose $a \neq 0$ to be a point that achieves the maximum of ϕ_k. We assume, without loss of generality, that $S_1(a)$ is non-negative (if $S_1(a) < 0$, consider $-a$ instead of a). There are two cases to consider:

The first case is that for all $i \in [n]$,

$$a_i \leq \frac{2k^{1/2}(S_1^2(a) + E_2(a))^{1/2}}{n}. \tag{9}$$

In this case we do not need the induction hypothesis and can in fact replace each a_i by its absolute value. Let $P \subseteq [n]$ be the set of $i \in [n]$ so that $a_i \geq 0$. Then by Equation (9),

$$\sum_{i \in P} |a_i| \leq 2k^{1/2}(S_1^2(a) + E_2(a))^{1/2}.$$

Note that

$$S_1(a) = \sum_{i \in P} |a_i| - \sum_{i \notin P} |a_i| \geq 0.$$

Hence

$$\sum_{i \notin P} |a_i| \leq \sum_{i \in P} |a_i| \leq 2k^{1/2}(S_1^2(a) + E_2(a))^{1/2}.$$

Overall we have

$$\sum_{i \in [n]} |a_i| \leq 4k^{1/2}(S_1^2(a) + E_2(a))^{1/2}.$$

We then bound

$$\begin{aligned}
|S_k(a_1, \ldots, a_n)| &\leq S_k(|a_1|, \ldots, |a_n|) \\
&\leq \left(\frac{e}{k}\right)^k \left(\sum_{i \in [n]} |a_i|\right)^k \quad \text{By the Maclaurin identities} \\
&\leq \left(\frac{4e}{\sqrt{k}}\right)^k (S_1^2(a) + E_2(a))^{k/2}.
\end{aligned}$$

[5]That is, for every $a \neq 0$ in $\mathbb{R}^n$ and $c \neq 0$ in $\mathbb{R}$, we have $\phi_k(ca) = \phi_k(a)$.

The second case is that there exists $i_0 \in [n]$ so that

$$a_{i_0} > \frac{2k^{1/2}(S_1^2(a) + E_2(a))^{1/2}}{n}. \tag{10}$$

In this case we use induction and Lagrange multipliers. For simplicity of notation, for a function F on $\mathbb{R}^n$ denote

$$F(-i) = F(a_1, a_2, \ldots, a_{i-1}, a_{i+1}, \ldots, a_n)$$

for $i \in [n]$. So, for every $\delta \in \mathbb{R}^n$ so that $\sum_i \delta_i = 0$ we have $\phi_k(a+\delta) \leq \phi_k(a)$. Hence[6], for all δ so that $\sum_i \delta_i = 0$,

$$\begin{aligned}
\phi_k(a) &\geq \frac{S_k^2(a+\delta)}{(S_1^2(a+\delta) + E_2(a+\delta))^k} \\
&\geq \frac{(S_k(a) + \sum_i \delta_i S_{k-1}(-i) + O(\delta^2))^2}{(S_1^2(a) + E_2(a) + 2\sum_i a_i\delta_i + O(\delta^2))^k} \\
&\geq \frac{S_k^2(a) + 2S_k(a)\sum_i \delta_i S_{k-1}(-i) + O(\delta^2)}{(S_1^2(a) + E_2(a))^k + 2k(S_1^2(a) + E_2(a))^{k-1}\sum_i a_i\delta_i + O(\delta^2)}.
\end{aligned}$$

Hence, for all δ close enough to zero so that $\sum_i \delta_i = 0$,

$$\frac{S_k^2(a)}{(S_1^2(a) + E_2(a))^k} \geq \frac{S_k^2(a) + 2S_k(a)\sum_i \delta_i S_{k-1}(-i) + O(\delta^2)}{(S_1^2(a) + E_2(a))^k + 2k(S_1^2(a) + E_2(a))^{k-1}\sum_i a_i\delta_i + O(\delta^2)},$$

or

$$\sum_i \delta_i \left(a_i S_k(a)k - (S_1^2(a) + E_2(a))S_{k-1}(-i)\right) \geq 0. \tag{11}$$

For the above inequality to hold for all such δ, it must be that there is λ so that for all $i \in [n]$,

$$a_i S_k(a)k - (S_1^2(a) + E_2(a))S_{k-1}(-i) = \lambda.$$

To see why this is true, set $\lambda_i = a_i S_k(a)k - (S_1^2(a) + E_2(a))S_{k-1}(-i)$. We now have $\lambda_1, \ldots, \lambda_n$ so that

$$\sum_i \lambda_i \delta_i \geq 0 \tag{12}$$

for every $\delta_1, \ldots, \delta_n$ of sufficiently small norm where $\sum_i \delta_i = 0$. We claim that this implies that in fact $\lambda_i = \lambda$ for every i. To see this, assume for contradiction that $\lambda_1 \neq \lambda_2$ and $|\lambda_1| > |\lambda_2|$. Set

$$\delta_1 = -\mu\lambda_1,\ \delta_2 = \mu\lambda_1,\ \delta_3 = \delta_4 = \ldots = \delta_n = 0$$

for $\mu > 0$ sufficiently small. It follows that $\sum_i \delta_i = 0$ and $\sum_i \lambda_i\delta_i = \mu(\lambda_1\lambda_2 - \lambda_1^2) < 0$ so Equation (12) is violated.

Sum over i to get

$$\lambda n = S_1(a)S_k(a)k - (S_1^2(a) + E_2(a))(n - (k-1))S_{k-1}(a).$$

[6]Here and below, $O(\delta^2)$ means of absolute value at most $C \cdot \|\delta\|_\infty$ for $C = C(n,k) \geq 0$.

Thus, for all $i \in [n]$,

$$\begin{aligned} a_i S_k(a)k - (S_1^2(a) + E_2(a))S_{k-1}(-i) \\ = \frac{1}{n}\left(S_1(a)S_k(a)k - (S_1^2(a) + E_2(a))(n-(k-1))S_{k-1}(a)\right), \end{aligned}$$

or

$$\begin{aligned} S_k(a)k\left(a_i - \frac{S_1(a)}{n}\right) \\ = (S_1^2(a) + E_2(a))(S_{k-1}(-i) - S_{k-1}(a)) + \frac{(k-1)}{n}(S_1^2(a) + E_2(a))S_{k-1}(a)). \end{aligned}$$

This specifically holds for i_0, so using (10) we have

$$\begin{aligned} &\left|S_k(a)k\frac{a_{i_0}}{2}\right| \\ &< \left|S_k(a)k\left(a_{i_0} - \frac{S_1(a)}{n}\right)\right| \\ &\leq \left|(S_1^2(a) + E_2(a))a_{i_0}S_{k-2}(-i_0)\right| + \left|\frac{(k-1)(S_1^2(a) + E_2(a))S_{k-1}(a)}{n}\right|, \end{aligned}$$

or

$$\begin{aligned} &|S_k(a)| \\ &\leq \left|\frac{2(S_1^2(a) + E_2(a))S_{k-2}(-i_0)}{k}\right| + \left|\frac{2(k-1)(S_1^2(a) + E_2(a))S_{k-1}(a)}{nka_{i_0}}\right| \\ &< \left|\frac{2(S_1^2(a) + E_2(a))S_{k-2}(-i_0)}{k}\right| + \left|\frac{(S_1^2(a) + E_2(a))^{1/2}S_{k-1}(a)}{k^{1/2}}\right|. \end{aligned} \tag{13}$$

To apply induction we need to bound $S_1^2(-i_0) + E_2(-i_0)$ from above. Since

$$\begin{aligned} S_1^2(a) + E_2(a) - S_1^2(-i_0) - E_2(-i_0) &= a_{i_0}^2 + 2a_{i_0}S_1(-i_0) + a_{i_0}^2 \\ &= 2a_{i_0}S_1(a) \geq 0. \end{aligned}$$

we have the bound

$$S_1^2(-i_0) + E_2(-i_0) \leq S_1^2(a) + E_2(a).$$

Finally, by induction and (13),

$$\begin{aligned} |S_k(a)| &\leq \frac{2(S_1^2(a) + E_2(a))}{k}\frac{(16e^2(S_1^2(-i_0) + E_2(-i_0)))^{(k-2)/2}}{(k-2)^{(k-2)/2}} \\ &\quad + \frac{(S_1^2(a) + E_2(a))^{1/2}}{k^{1/2}}\frac{(16e^2(S_1^2(a) + E_2(a)))^{(k-1)/2}}{(k-1)^{(k-1)/2}} \\ &\leq \frac{(16e^2(S_1^2(a) + E_2(a)))^{k/2}}{k^{k/2}}\left(\frac{2}{16e^2\left(1-\frac{2}{k}\right)^{(k-2)/2}} + \frac{1}{4e\left(1-\frac{1}{k}\right)^{(k-1)/2}}\right) \\ &< \frac{(16e^2(S_1^2(a) + E_2(a)))^{k/2}}{k^{k/2}}. \end{aligned}$$

□

Proof of Theorem 3. The proof is by reduction to Theorem 2. Assume $a_1, \ldots, a_m$ are nonzero and $a_{m+1}, \ldots, a_n$ are zero. Denote $a' = (a_1, \ldots, a_m)$ and notice that for all[7] $k \in [n]$,

$$S_k(a) = S_k(a').$$

Write

$$p(\xi) = \prod_{i \in [m]} (\xi a_i + 1) = \sum_{k=0}^{m} \xi^k S_k(a).$$

Derive k times to get

$$p^{(k)}(\xi) = S_k(a)k! \left(\binom{m}{k} \frac{S_m(a)}{S_k(a)} \xi^{m-k} + \binom{m-1}{k} \frac{S_{m-1}(a)}{S_k(a)} \xi^{m-k-1} + \ldots \right.$$
$$\left. \ldots + \binom{k+1}{k} \frac{S_{k+1}(a)}{S_k(a)} \xi + 1 \right).$$

Since p has m real roots, $p^{(k)}$ has $m-k$ real roots. Since $p^{(k)}(0) \neq 0$, there is $b \in \mathbb{R}^{m-k}$ so that

$$p^{(k)}(\xi) = S_k(a)k! \prod_{i \in [m-k]} (\xi b_i + 1).$$

For all $h \in [m-k]$,

$$S_h(b) = \binom{k+h}{k} \frac{S_{k+h}(a)}{S_k(a)}.$$

By assumption,

$$|S_1(b)| \leq C \quad \text{and} \quad |S_2(b)| \leq C^2.$$

Theorem 2 implies

$$|S_h(b)| = \left| \binom{k+h}{k} \frac{S_{k+h}(a)}{S_k(a)} \right| \leq \frac{(6eC)^h}{h^{h/2}}.$$

□

3 Tail bounds under limited independence

In this section we work with the following setup: Let $X = (X_1, \ldots, X_n)$ be a vector of real valued random variables so that $\mathbb{E}[X_i] = 0$ for all $i \in [n]$. Denote $\sigma_i^2 = \mathsf{Var}[X_i]$ and

$$\sigma^2 = \sum_{i \in [n]} \sigma_i^2.$$

The goal is proving a tail bound on the behaviour of the symmetric functions under limited independence.

We start by obtaining tail estimates, under full independence. Let $\mathcal{U}$ denote the distribution over $X = (X_1, \ldots, X_n)$ where $X_1, \ldots, X_n$ are independent.

Lemma 9. $\mathbb{E}_{X \in \mathcal{U}}[S_\ell^2(X)] \leq \frac{\sigma^{2\ell}}{\ell!}$.

[7] For $k > m$ we have $S_k(a) = 0$ so there is nothing to prove.

Proof. Since the expectation of X_i is zero for all $i \in [n]$,

$$\begin{aligned}\mathbb{E}[S_\ell^2(X)] &= \sum_{T,T' \subset [n]:|T|=|T'|=\ell} \mathbb{E}\left[\prod_{t\in T} X_t \prod_{t'\in T'} X_{t'}\right] \\ &= \sum_{T\subset[n]:|T|=\ell} \mathbb{E}\left[\prod_{t\in T} X_t^2\right] = \sum_{T\subset[n]:|T|=\ell} \prod_{t\in T} \sigma_t^2 \\ &\le \frac{1}{\ell!}\left(\sum_{i\in[n]} \sigma_i^2\right)^\ell = \frac{\sigma^{2\ell}}{\ell!}.\end{aligned}$$

□

Corollary 10. *For $t > 0$ and $\ell \in [n]$, by Markov's inequality,*

$$\Pr_{X\in\mathcal{U}}\left[|S_\ell(X)| \ge \left(\frac{e^{1/2}t\sigma}{\ell^{1/2}}\right)^\ell \ge \frac{(t\sigma)^\ell}{\sqrt{\ell!}}\right] \le \frac{1}{t^{2\ell}}. \tag{14}$$

If $2e^{1/2}t\sigma \le k^{1/2}$ then by the union bound

$$\Pr_{X\in\mathcal{U}}\left[\sum_{\ell=k}^{n} |S_\ell(X)| \ge 2\left(\frac{e^{1/2}t\sigma}{k^{1/2}}\right)^k\right] \le \frac{1}{t^{2k} - t^{2(k-1)}}. \tag{15}$$

We now consider limited independence.

Lemma 11. *Let $\mathcal{D}$ denote a distribution over $X = (X_1, \ldots, X_n)$ where $X_1, \ldots, X_n$ are $(2k+2)$-wise independent. Let $t \ge 1$. Except with $\mathcal{D}$-probability at most $2t^{-2k}$, the following bounds hold for all $\ell \in \{k, \ldots, n\}$:*

$$|S_\ell(X)| \le (6et\sigma)^\ell \left(\frac{k}{\ell}\right)^{\ell/2}. \tag{16}$$

Proof. In the following the underlying probability distribution over X is $\mathcal{D}$. By Lemma 9, for $i \in \{k, k+1\}$,

$$\mathbb{E}[S_i^2(X)] \le \frac{\sigma^{2i}}{i!}.$$

Hence by Markov's inequality,

$$\Pr\left[|S_i(X)| \ge \frac{(t\sigma)^i}{\sqrt{i!}}\right] \le t^{-2i}.$$

From now on, condition on the event that

$$|S_k(X)| \le \frac{(t\sigma)^k}{\sqrt{k!}} \text{ and } |S_{k+1}(X)| \le \frac{(t\sigma)^{k+1}}{\sqrt{(k+1)!}}, \tag{17}$$

which occurs with probability at least $1 - 2t^{-2k}$. Fix $x = (x_1, \ldots, x_n)$ such that Equation (17) holds.

We claim that there must exist $k_0 \in \{0, \ldots, k-1\}$ for which the following bounds hold:

$$|S_{k_0}(x)| \geq \frac{(t\sigma)^{k_0}}{\sqrt{k_0!}}, \tag{18}$$

$$|S_{k_0+1}(x)| \leq \frac{(t\sigma)^{k_0+1}}{\sqrt{(k_0+1)!}}, \tag{19}$$

$$|S_{k_0+2}(x)| \leq \frac{(t\sigma)^{k_0+2}}{\sqrt{(k_0+2)!}}. \tag{20}$$

To see this, mark point $j \in \{0, \ldots, k+1\}$ as *high* if

$$|S_j(x)| \geq \frac{(t\sigma)^j}{\sqrt{j!}}$$

and *low* if

$$|S_j(x)| \leq \frac{(t\sigma)^j}{\sqrt{j!}}.$$

A point is marked both high and low if equality holds. Observe that 0 is marked high (and low) since $S_0(x) = 1$ and k and $k+1$ are marked low by Equation (17). This implies the existence of a triple k_0, k_0+1, k_0+2 where the first point is high and the next two are low.

Let $\gamma > 0$ be the smallest number so that the following inequalities hold:

$$|S_{k_0+1}(x)| \leq |S_{k_0}(x)| \frac{\gamma}{\sqrt{k_0+1}}, \tag{21}$$

$$|S_{k_0+2}(x)| \leq |S_{k_0}(x)| \frac{\gamma^2}{\sqrt{(k_0+1)(k_0+2)}}. \tag{22}$$

By definition, one of Equations (21) and (22) holds with equality so

$$|S_{k_0}(x)| = \max\left\{ \frac{|S_{k_0+1}(x)|\sqrt{k_0+1}}{\gamma}, \frac{|S_{k_0+2}(x)|\sqrt{(k_0+1)(k_0+2)}}{\gamma^2} \right\}.$$

Observe further that $\gamma \leq t\sigma$ by Equations (18), (19) and (20). Combining this with the bounds in Equations (19) and (20)

$$|S_{k_0}(x)| \leq \max\left\{ \frac{(t\sigma)^{k_0+1}}{\gamma\sqrt{k_0!}}, \frac{(t\sigma)^{k_0+2}}{\gamma^2\sqrt{k_0!}} \right\} = \frac{(t\sigma)^{k_0+2}}{\gamma^2\sqrt{k_0!}}. \tag{23}$$

Equations (21) and (22) let us apply Theorem 3 with $C = \gamma\sqrt{k_0+1}$ and $h \geq 3$ to get

$$\left| \frac{S_{k_0+h}(x)}{S_{k_0}(x)} \right| \leq (6e\gamma)^h \frac{(k_0+1)^{h/2}}{h^{h/2}\binom{k_0+h}{k_0}}.$$

Bounding $|S_{k_0}|$ by Equation (23), we get

$$|S_{k_0+h}(x)| \le (6e\gamma)^h \frac{(k_0+1)^{h/2}}{h^{h/2}\binom{k_0+h}{k_0}} \frac{(t\sigma)^{k_0+2}}{\gamma^2\sqrt{k_0!}} \le (6et\sigma)^{k_0+h} \frac{(k_0+1)^{h/2}}{h^{h/2}\sqrt{k_0!}\binom{k_0+h}{h}}.$$

Since

$$\binom{k_0+h}{h} \ge \max\left\{\left(\frac{k_0+h}{k_0}\right)^{k_0}, \left(\frac{k_0+h}{h}\right)^{h}\right\} \ge \frac{(k_0+h)^{(k_0+h)/2}}{k_0^{k_0/2}h^{h/2}},$$

we have

$$\frac{(k_0+1)^{h/2}}{h^{h/2}\sqrt{k_0!}\binom{k_0+h}{h}} \le \left(\frac{k_0+1}{h}\right)^{h/2} \frac{k_0^{k_0/2}h^{h/2}}{(k_0+h)^{(k_0+h)/2}} \le \left(\frac{k_0+1}{k_0+h}\right)^{(k_0+h)/2}.$$

Therefore, denoting $\ell = k_0 + h$, since $k_0 + 1 \le k$,

$$|S_\ell(x)| \le (6et\sigma)^\ell \left(\frac{k}{\ell}\right)^{\ell/2}.$$

□

Proof of Theorem 4. As in Lemma 11, fix $x = (x_1, \ldots, x_n)$ such that Equation (17) holds (the random vector X has this property with $\mathcal{D}$-probability at least $1 - 2t^{-2k}$). By the proof of lemma, since by assumption $6et\sigma < 1/2$,

$$\sum_{\ell=k}^{n} |S_\ell(x)| \le \frac{(t\sigma)^k}{k!} + \frac{(t\sigma)^{k+1}}{\sqrt{(k+1)!}} + \sum_{\ell=k+2}^{n} (6et\sigma)^\ell \left(\frac{k}{\ell}\right)^{\ell/2} \le 2(6et\sigma)^k. \tag{24}$$

□

3.1 On the tightness of the tail bounds

We conclude by showing that $(2k+2)$-wise independence is insufficient to fool $|S_\ell|$ for $\ell > 2k+2$ in expectation. We use a modification of a simple proof due to Noga Alon of the $\Omega(n^{k/2})$ lower bound on the support size of a k-wise independent distribution on $\{-1, 1\}^n$, which was communicated to us by Raghu Meka.

For this section, let $X_1, \ldots, X_n$ be so that each X_i is uniform over $\{-1, 1\}$. Thus $\sigma^2 = \sum_i \mathsf{Var}[X_i] = n$. By Lemma 9, we have

$$\mathbb{E}_{X\in\mathcal{U}}[|S_\ell(X)|] \le \left(\mathbb{E}_{X\in\mathcal{U}}[S_\ell^2(X)]\right)^{1/2} \le \frac{n^{\ell/2}}{\sqrt{\ell!}}. \tag{25}$$

In contrast we have the following:

Lemma 12. *There is a $(2k+2)$-wise independent distribution on $X = (X_1, X_2, \ldots, X_n)$ in $\{-1, 1\}^n$ such that for every $\ell \in [n]$,*

$$\Pr_{X\in\mathcal{D}}\left[|S_\ell(X)| \ge \binom{n}{\ell}\right] \ge \frac{1}{3n^{k+1}}.$$

Specifically,

$$\mathbb{E}_{X\in\mathcal{D}}[|S_\ell(X)|] \geq \frac{\binom{n}{\ell}}{3n^{k+1}}. \tag{26}$$

Proof. Let $\mathcal{D}$ be a $(2k+2)$-wise independent distribution on $\{-1,1\}^n$ that is uniform over a set D of size $2(n+1)^{k+1} \leq 3n^{k+1}$. Such distributions are known to exist [ABI86]. Further, by translating the support by some fixed vector if needed, we may assume that $(1,1,\ldots,1) \in D$. It is easy to see that every such translate also induces a $(2k+2)$-wise independent distribution. The claim holds since $S_\ell(1,\ldots,1) = \binom{n}{\ell}$. □

When e.g. $k = O(\log n)$, which is often the case of interest, for $2k+3 \leq \ell \leq n-(2k+3)$, the RHS of (26) is much larger than the bound guaranteed by Equation (25). The tail bound provided by Lemma 11 can not therefore be extended to a satisfactory bound on the expectation. Furthermore, applying Lemma 11 with

$$t = \frac{1}{6e}\sqrt{\frac{n}{\ell k}}$$

implies that for any $(2k+2)$-wise independent distribution,

$$\Pr\left[|S_\ell(X)| \geq \binom{n}{\ell}\right] \leq \Pr\left[|S_\ell(X)| \geq (6et\sqrt{n})^\ell \left(\frac{k}{\ell}\right)^{\ell/2}\right] \leq 2\left(\frac{36e^2k\ell}{n}\right)^k.$$

When $k\ell = o(n)$, this is at most $O(n^{-k+o(1)})$. Comparing this to the bound given in Lemma 12, we see that the bound provided by Lemma 11 is nearly tight.

4 Limited independence fools products of bounded variables

In this section we work with the following setup. We have n random variables $X_1,\ldots,X_n$ each distributed in the interval $[-1,1]$. Let μ_i and σ_i^2 denote the mean and variance of X_i, and let $\sigma^2 = \sum_{i=1}^n \sigma_i^2$. We will typically use $\mathcal{U}$ to denote the distribution where the X_is are fully independent, and $\mathcal{D}$ to denote distributions with limited independence.

Theorem 13. *There exist constants $c, c' > 0$ such that under any ck-wise independent distribution $\mathcal{D}$,*

$$\left|\mathbb{E}_{\mathcal{D}}[\prod_{i=1}^n X_i] - \prod_{i=1}^n \mu_i\right| \leq (c'\sigma)^k. \tag{27}$$

Proof. Define $H \subset [n]$ to be the set of indices such that $|\mu_i| \leq \sqrt{\sigma}$. Note that if $H \geq 2k$, then we are done since if $c \geq 2$, then

$$\left|\mathbb{E}_{\mathcal{D}}[\prod_{i\in H} X_i]\right| = \prod_{i\in H}|\mu_i| \leq \sqrt{\sigma}^{2k} \leq \sigma^k$$

Further, since the variables are bounded in $[-1,1]$, we have

$$\left|\prod_{i\in n} X_i\right| \leq \left|\prod_{i\in H} X_i\right|$$

hence

$$\left|\mathbb{E}_{\mathcal{D}}[\prod_{i\in[n]} X_i]\right| \leq \left|\mathbb{E}_{\mathcal{D}}[\prod_{i\in H} X_i]\right| \leq \sigma^k.$$

The same bound also holds under $\mathcal{U}$, hence

$$\left|\mathbb{E}_{\mathcal{D}}[\prod_{i\in[n]} X_i] - \mathbb{E}_{\mathcal{U}}[\prod_{i\in[n]} X_i]\right| \leq 2\sigma^k.$$

So now assume that $|H| \leq 2k$. Let $T = H \setminus [n]$. Even after conditioning on the outcome of variables in H, the resulting distribution on T is $(c-2)k = c''k$-wise independent. Since the product of variables in H has absolute value at most 1, it suffices to show that for a $c''k$-wise independent distribution $\mathcal{D}$,

$$\left|\mathbb{E}_{\mathcal{D}}[\prod_{i\in T} X_i] - \mathbb{E}_{\mathcal{U}}[\prod_{i\in T} X_i]\right| \leq 2\sigma^k.$$

For ease of notation, we shall assume that $T = [m]$ for some $m \leq n$. We may assume that $m > c''k$ else there is nothing to prove.

Let us write $X_i = \mu_i(1+Z_i)$, so that Z_i has mean 0 and variance σ_i^2/μ_i^2. We write

$$\prod_{i\in[m]} X_i = \prod_{i\in[m]} \mu_i(1+Z_i) = \prod_{i\in[m]} \mu_i \left(\sum_{\ell\leq m} S_\ell(Z_1,\ldots,Z_m)\right)$$

. Let us define the functions

$$P(Z) = \prod_{i\in[m]} \mu_i(1+Z_i)$$
$$P'(Z) = \prod_{i\in[m]} \sum_{\ell\leq 4k} S_\ell(Z_1,\ldots,Z_m).$$

We will prove the following claim.

Claim 14. *For a $c''k$-wise independent distribution $\mathcal{D}$,*

$$\left|\mathbb{E}_{\mathcal{D}}[P(Z)] - \mathbb{E}_{\mathcal{D}}[P'(Z)]\right| \leq (c'\sigma)^k/2.$$

We first show how to finish the proof of Theorem 13 with this claim. We have

$$\begin{aligned}|\mathbb{E}_{\mathcal{D}}[P(Z)] - \mathbb{E}_{\mathcal{U}}[P(Z)]| \leq \left|\mathbb{E}_{\mathcal{D}}[P(Z)] - \mathbb{E}_{\mathcal{D}}[P'(Z)]\right| + \left|\mathbb{E}_{\mathcal{U}}[P(Z)] - \mathbb{E}_{\mathcal{U}}[P'(Z)]\right| \\ + \left|\mathbb{E}_{\mathcal{D}}[P'(Z)] - \mathbb{E}_{\mathcal{U}}[P'(Z)]\right|\end{aligned}$$

The first two are bounded by $(c'\sigma^k)/2$ by the claim, and the last is 0 since $c''k$-wise independence fools degree $4k$ polynomials for $c'' > 4$.

Proof of Claim 14. Recall that the X_is for $i \in [m]$ have expectation μ_i where $|\mu_i| \geq \sqrt{\sigma}$. We let $X_i = \mu_i(1 + Z_i)$, where Z_i has mean 0 and variance $\bar{\sigma}_i^2$ where

$$\bar{\sigma}_i^2 = \frac{\sigma_i^2}{\mu_i^2} \leq \frac{\sigma_i^2}{\sigma}.$$

Hence the total variance of the Z_is can be bounded by

$$\bar{\sigma}^2 \leq \sum_{i \in T} \frac{\sigma_i^2}{\sigma} \leq \sigma.$$

Writing $Z = (Z_1, \ldots, Z_m)$ we have

$$P(Z) - P'(Z) = \sum_{\ell=4k+1}^{m} |S_\ell(Z)|.$$

Let G denote the event that $|P(Z) - P'(Z)| \leq 2(6e\sqrt{\bar{\sigma}})^{4k}$. Letting $t = 1/\sqrt{\bar{\sigma}}$ and applying Theorem 4, for $c'' > 8k + 2$

$$\mathbb{E}_{\mathcal{D}}[\mathbb{1}(\neg G)] \leq 2t^{-8k} = 2\bar{\sigma}^{4k}. \tag{28}$$

Since $\mathbb{E}[Z_i] = 0$ for all i it follows that under $c''k$-wise independence,

$$\mathbb{E}[P_k(Z_1, \ldots, Z_n)^2] \leq \sum_{i=0}^{4k} \mathbb{E}[S_i(Z_1, \ldots, Z_n)^2] \leq \sum_{i=0}^{4k} \frac{\bar{\sigma}^{2i}}{i!} \leq 2. \tag{29}$$

We now write

$$\mathbb{E}[P(Z) - P'(Z)] = \mathbb{E}[(P(Z) - P'(Z))\mathbb{1}(G)] + \mathbb{E}[(P(Z) - P'(Z))\mathbb{1}(\neg G)].$$

Equation (37) implies

$$|\mathbb{E}[(P(Z) - P'(Z))\mathbb{1}(G)]| \leq 2(6e\sqrt{\bar{\sigma}})^{4k}.$$

For the second term,

$$|\mathbb{E}[(P(Z) - P'(Z))\mathbb{1}(\neg G)]| \leq |\mathbb{E}[P(Z)\mathbb{1}(\neg G)]| + |\mathbb{E}[P'(Z)\mathbb{1}(\neg G)]|$$

Note that $0 \leq P(Z) \leq 1$. Also note that $\mathbb{E}[P_k(Z)^2] \leq 2$ by Equation (29). So we can bound the RHS using Holder's inequalities by

$$|\mathbb{E}[\mathbb{1}(\neg G)]| + |\mathbb{E}[P_k(A)^2]^{1/2} \cdot \mathbb{E}[\mathbb{1}(\neg G)]^{1/2}| \leq \bar{\sigma}^{4k} + \sqrt{2}\bar{\sigma}^{2k} \leq 2\bar{\sigma}^{2k}.$$

Hence overall we have

$$\mathbb{E}[P(Z) - P'(Z)] = 2(6e\sqrt{\bar{\sigma}})^{4k} + 2\bar{\sigma}^{2k} \leq (c'\sigma)^k/2$$

□

□

5 Analyzing the [GMR$^+$12] generator

Gopalan et al. [GMR$^+$12] proposed and analyzed a PRG for combinatorial rectangles, which we denote by $\mathcal{G}_{\mathcal{MR}}$. In this section, we provide a different analysis of their construction, which is based on our results concerning the symmetric polynomials. Our analysis is simpler and follows the intuition that products of low variance events are easy to fool using limited independence. It also improves one their seedlength in the dependence on n, δ (see the discussion following Theorem 7).

Let $\mathcal{U}$ denote the uniform distribution on $[m]^n$, and let $\mathcal{D}$ be a distribution on $[m]^n$. For $x \in [m]^n$ and $K \subseteq [n]$, let $x_K = (x_i : i \in K)$. We sometimes abuse notation and write x_K instead of the probability distribution of x_K. We denote by d_{TV} the total variation distance.

Definition 15. A distribution $\mathcal{D}$ on $[m]^n$ is (k, ε)-wise independent if for every $K \subseteq [n]$ of size k, and $x \in \mathcal{D}, y \in \mathcal{U}$, we have $d_{TV}(x_K, y_K) \leq \varepsilon$.

Such distributions can be generated using seed length $O(\log\log(n) + k\log(m) + \log(1/\varepsilon))$ when m is a power of 2 using standard constructions [NN93]. We can also assume that every co-ordinate is uniformly random in $[m]$. See the appendix for details.

(by adding the string $(a, a, \ldots, a)$ modulo m, where $a \in [m]$ is uniformly random).

Being (k, ε)-wise independent is equivalent to saying that for every $K \subseteq [n]$ of size k and every $f : [m]^k \to \{0, 1\}$,

$$|\mathbb{E}_{x\in\mathcal{D}}[f(x_K)] - \mathbb{E}_{y\in\mathcal{U}}[f(y_K)]| \leq \varepsilon.$$

The following more general property holds. Let P be a real linear combination of combinatorial rectangles,

$$P = \sum_{S\subseteq[n]} c_S f_S,$$

where $f_S(x) = \prod_{i\in S} f_i(x_i)$. Let $\mathrm{L}_1(P) = \sum_S |c_S|$. The degree of P is the maximum size of S for which $c_S \neq 0$. It follows that if $\mathcal{D}$ is (k, ε)-wise independent and P has degree at most k then

$$|\mathbb{E}_{x\in\mathcal{D}}[P(x)] - \mathbb{E}_{x\in\mathcal{U}}[P(x)]| \leq \mathrm{L}_1(P)\varepsilon.$$

5.1 The generator

We use an alternate view of $\mathcal{G}_{\mathcal{MR}}$ as a collection of hash functions $g : [n] \to [m]$. The generator $\mathcal{G}_{\mathcal{MR}}$ is based on iterative applications of an *alphabet increasing* step. The first alphabet m_0 is chosen to be large enough, and at each step $t > 1$ the size of the alphabet m_t is squared $m_t = m_{t-1}^2$. There is a constant $C > 0$ so that the following holds. Denote by δ the error parameter of the generator. Let $T \leq C\log\log(m)$ be the first integer so that $m_T \geq m$. Let $\delta' = \delta/T$.

1. **Base Case:** Let $m_0 \geq C\log(1/\delta)$ be a power of 2. Sample $g_0 : [n] \to [m_0]$ using a (k_0, ε_0)-wise independent distribution on $[m_0]^n$ with

$$k_0 = C\log(1/\delta'),\ \varepsilon_0 = \delta' \cdot m_0^{-Ck_0}. \tag{30}$$

   This requires seed length $O(\log\log(n) + \log(\log\log(m)/\delta)\log\log(\log\log(m)/\delta))$.

2. **Squaring the alphabet:** Pick $g'_t : [m_{t-1}] \times [n] \to [m_t]$ using a (k_t, ε_t)-wise independent distribution over $[m_t]^{m_{t-1} \times n}$ with

$$k_t = \max\left\{C\frac{\log(1/\delta')}{\log(m_t)}, 2\right\},\ \varepsilon_t \leq m_t^{-Ck_t}.$$

Define a hash function $g_t : [n] \to [m_t]$ as

$$g_t(i) = g'_t(g_{t-1}(i), i).$$

This requires seed length $O(\log\log(n) + \log(m_t) + \log(\log\log(m)/\delta))$.

5.2 Analyzing the generator

We first analyze the base case using the inclusion-exclusion approach of [EGL+98]. We need to extend their analysis to the setting where the co-ordinates are only approximately k-wise independent.

Lemma 16. *Let $\mathcal{D}$ be a (k, ε)-wise independent distribution on $[m]^n$ with k odd. Then,*

$$\left|\Pr_{g\in\mathcal{D}}[\forall\ i \in [n]\ g(i) \in S_i] - \Pr_{h\in\mathcal{U}}[\forall\ i \in [n]\ h(i) \in S_i]\right| \leq \varepsilon m^k + \exp(-\Omega(k)).$$

Proof. Let $p_i = |S_i|/m$, and $q_i = 1 - p_i$. Observe that

$$\Pr_h[\forall\ i \in [n]\ h(i) \in S_i] = \prod_{i=1}^{n} p_i = \prod_{i=1}^{n}(1 - q_i) \leq \exp\left(-\sum_{i=1}^{n} q_i\right). \tag{31}$$

We consider two cases based on $\sum_i q_i$.

Case 1: When $\sum_i q_i \leq k/(2e)$. Since every non-zero q_i is at least $1/m$, there can be at most $mk/(2e)$ indices i so that $q_i > 0$. For i so that $q_i = 0$, we have $S_i = [m]$, so we can drop such indices and assume $n \leq mk/(2e)$. By Bonferroni inequality, since k is odd,

$$\left|\Pr_g[\forall\ i \in [n]\ g(i) \in S_i] - \sum_{j=0}^{k-1}(-1)^j \sum_{J\subseteq[n]:|J|=j} \Pr_g[\forall\ i \in J\ g(i) \notin S_i]\right|$$
$$\leq \sum_{J\subseteq[n]:|J|=k} \Pr_g[\forall\ i \in J\ g(i) \notin S_i].$$

A similar bound holds for h. The (k, ε)-wise independence thus implies

$$\left|\Pr_g[\forall\ i \in [n]\ g(i) \in S_i] - \Pr_h[\forall\ i \in [n]\ h(i) \in S_i]\right|$$
$$\leq \varepsilon(en/k)^k + 2\sum_{J\subseteq[n]:|J|=k} \Pr_h[\forall\ i \in J\ h(i) \notin S_i].$$

The second term is twice $S_k(q_1, \ldots, q_n)$, which we can bound by Maclaurin's identity as

$$S_k(q_1, \ldots, q_n) \leq (e/k)^k \left(\sum_{i=1}^{n} q_i\right)^k \leq 2^{-k}.$$

Finally, since $n \le mk/(2e)$,

$$\left|\Pr_g[\forall\ i \in [n]\ g(i) \in S_i] - \Pr_h[\forall\ i \in [n]\ h(i) \in S_i] - S_i]\right| \le \varepsilon(m/2)^k + 2^{-k+1}.$$

Case 2: When $\sum_i q_i > k/2e$. Once again, we drop indices i so that $q_i = 0$. Consider the largest n' such that

$$k/2e - 1 \le \sum_{i=1}^{n'} q_i \le k/2e.$$

Repeating the argument from Case 1 for this n',

$$\left|\Pr_g[\forall\ i \in [n']\ g(i) \in S_i] - \Pr_h[\forall\ i \in [n']\ h(i) \in S_i]\right| \le \varepsilon(m/2)^k + 2^{-k+1}.$$

Similarly to Equation (31),

$$\Pr_h[\forall\ i \in [n']\ h(i) \in S_i] \le e^{-k/2e+1}.$$

Since

$$\Pr_g[\forall\ i \in [n]\ g(i) \in S_i] \le \Pr_g[\forall\ i \in [n']\ g(i) \in S_i],$$

we have

$$\left|\Pr_g[\forall\ i \in [n]\ g(i) \in S_i] \le \Pr_h[\forall\ i \in [n]\ h(i) \in S_i]\right| \le \varepsilon(m/2)^k + \exp(-\Omega(k)).$$

□

To analyze the iterative steps, we use the following lemma:

Lemma 17. *There is $C > 0$ so that the following holds for $\delta > 0$ small enough. Assume*

$$k > 1,\ \ell \ge \log(1/\delta),\ \ell \ge k,\ \ell^{-k} \le \delta^C,\ \varepsilon \le (m\ell)^{-Ck}.$$

Let $\mathcal{D}$ be a (k, ε)-wise independent distribution on $g' : [\ell] \times [n] \to [m]$ so that for every $(a, i) \in [\ell] \times [n]$ the distribution of $g'(a, i)$ is uniform on $[m]$. Then,

$$\left|\Pr_{g' \in \mathcal{D}, x \in [\ell]^n}[\forall\ i \in [n]\ g'(x_i, i) \in S_i] - \Pr_{h \in [m]^n}[\forall\ i \in [n]\ h(i) \in S_i]\right| \le \delta.$$

Proof. Given g', x, let $g : [n] \to [m]$ be defined by $g(i) = g'(x_i, i)$. We can similarly pick h in two steps: pick $h' : [\ell] \times [n] \to [m]$ uniformly at random, pick $x \in [\ell]^n$ independently and uniformly at random, and then let $h(i) = h'(x_i, i)$.

For every $i \in [n]$, since each x_i is uniform over $[\ell]$, for every fixed g', we have

$$\Pr_x[g(i) \in S_i] = \frac{1}{\ell}\sum_{a=1}^{\ell} \mathbb{1}(g'(a, i) \in S_i).$$

So, for every fixed g',

$$\Pr_x[\forall\ i \in n\ g(i) \in S_i] = \prod_{i=1}^{n} \Pr_x[g(i) \in S_i] = \prod_{i=1}^{n} \frac{1}{\ell} \sum_{a=1}^{\ell} \mathbb{1}(g'(a,i) \in S_i). \tag{32}$$

A similar equation holds for h.

Let $p_i = |S_i|/m$ and $q_i = 1 - p_i$. Partition $[n]$ into a head $H = \{i : p_i < \ell^{-0.1}\}$ and a tail $T = \{i : p_i \geq \ell^{-0.1}\}$. Standard arguments (see e.g. [GMR$^+$12, Theorem 4.1]) imply that if (k,ε)-wise independence fools both $\forall\ i \in H\ g(i) \in S_i$ and $\forall\ i \in T\ g(i) \in S_i$ with error δ then $(O(k), \varepsilon^{O(1)})$-wise independence fools their intersection with error $O(\delta)$. So it suffices to consider each of them separately.

Fooling the Head: If $|H| \leq k$,

$$\Pr_x[\forall\ i \in H\ g(i) \in S_i] = \prod_{i \in H} \frac{1}{\ell} \sum_{a=1}^{\ell} \mathbb{1}(g'(a,i) \in S_i)$$

is a degree k polynomial with L_1-norm bounded by 1. Hence,

$$\left| \mathbb{E}_{g'}[\Pr_x[\forall\ i \in H\ g(i) \in S_i]] - \mathbb{E}_{h'}[\Pr_x[\forall\ i \in H\ h(i) \in S_i]] \right| \leq \varepsilon \leq \delta. \tag{33}$$

If $|H| \geq k$, we show that the probabilities are small which means that they are close. Indeed, let H' be the first k indices in H. First,

$$\Pr_h[\forall\ i \in H'\ h(i) \in S_i] = \prod_{i \in H'} \Pr[h(i) \in S_i] \leq \ell^{-0.1k} \leq \delta.$$

Second, Equation (33) implies

$$\Pr_g[\forall\ i \in H'\ g(i) \in S_i] \leq \ell^{-0.1k} + \varepsilon \leq \delta.$$

Fooling the Tail: We may assume that $q_i \geq 1/m$ and $p_i > 0$ for all $i \in T$, since otherwise S_i is trivial and we can drop such an index. As in the proof of Lemma 16, by restricting to a subset if necessary, we can also assume that

$$\sum_{i \in T} q_i \leq C \log(1/\delta). \tag{34}$$

For simplicity of notation, we denote $|T|$ by n. Therefore, $n \leq Cm \log(1/\delta)$.

Let

$$Y(a,i) = \mathbb{1}(g'(a,i)) - p_i.$$

Since $g'(a,i)$ is uniform over S_i,

$$\mathbb{E}[Y(a,i)] = 0,\ \mathsf{Var}[Y(a,i)] = q_i p_i.$$

Write

$$\Pr_x[\forall\ i \in T\ g(i) \in S_i] = \prod_{i \in T} \left(p_i + \frac{1}{\ell} \sum_{a=1}^{\ell} Y(a,i) \right).$$

Define new random variables

$$A_i = \frac{1}{\ell p_i} \sum_{a=1}^{\ell} Y(a,i).$$

so that

$$\Pr_x[\forall\ i \in T\ g(i) \in S_i] = \prod_{i=1}^{n} p_i(1+A_i) = \prod_{i=1}^{n} p_i \cdot \left(\sum_{i=0}^{n} S_i(A_1, \ldots, A_n) \right). \tag{35}$$

For $k \leq n$, define

$$P_k(A) = \prod_{i=1}^{n} p_i \cdot \left(\sum_{i=0}^{k} S_i(A_1, \ldots, A_n) \right).$$

We will show that $P_k(A)$ is a good approximation to $P_n(A)$ under $(O(k), \varepsilon^{O(1)})$-wise independence, hence under both $\mathcal{D}$ and $\mathcal{U}$.

Claim 18. *Both $|\mathbb{E}_{\mathcal{D}}[P_n(A) - P_k(A)]|$ and $|\mathbb{E}_{\mathcal{U}}[P_n(A) - P_k(A)]|$ are at most $O(\ell^{-0.2k})$.*

The claim completes the proof:

$$\begin{aligned}|\mathbb{E}_{\mathcal{D}}[P_n(A)] - \mathbb{E}_{\mathcal{U}}[P_n(A)]| \leq |\mathbb{E}_{\mathcal{D}}[P_n(A)] - \mathbb{E}_{\mathcal{D}}[P_k(A)]| + |\mathbb{E}_{\mathcal{D}}[P_k(A)] - \mathbb{E}_{\mathcal{U}}[P_k(A)]| \\ + |\mathbb{E}_{\mathcal{U}}[P_n(A)] - \mathbb{E}_{\mathcal{U}}[P_k(A)]|.\end{aligned}$$

Bound the first and third terms by $O(\ell^{-0.2k})$ using the claim. Bound the second term as follows. Since $k > 1$, for all i,

$$\mathsf{Var}[A_i] = \frac{1}{\ell^2 p_i^2} \sum_{a=1}^{\ell} \mathsf{Var}[Y(a,i)] = \frac{q_i}{\ell p_i} \leq \frac{q_i}{\ell^{0.9}},$$

$$\mathrm{L}_1(A_i) \leq \frac{1}{\ell p_i} \sum_{a=1}^{\ell} \mathrm{L}_1(Y(a,i)) \leq \frac{2}{p_i} \leq \ell.$$

Plugging in the bounds from Equations (34):

$$\sum_{i=1}^{n} \mathsf{Var}[A_i] \leq \frac{C \log(1/\delta)}{\ell^{0.9}} \leq \frac{1}{\ell^{0.6}},$$

$$\sum_{i=1}^{n} \mathrm{L}_1(A_i) \leq Cm \log(1/\delta)\ell \leq m\ell^{O(1)},$$

$$\mathrm{L}_1(S_k(A_1, \ldots, A_n)) \leq \left(\sum_{i=1}^{n} \mathrm{L}_1(A_i) \right)^k \leq m^k \ell^{O(k)}.$$

Thus,

$$|\mathbb{E}_{\mathcal{D}}[P_k(A)] - \mathbb{E}_{\mathcal{U}}[P_k(A)]| \leq \varepsilon \mathrm{L}_1(P_k) \leq \varepsilon \ell^{O(k)} = O(\ell^{-k}).$$

Overall,

$$\begin{aligned}\left| \Pr_g[\forall\ i \in T\ g(i) \in S_i] - \Pr_h[\forall\ i \in T\ h(i) \in S_i] \right| \\ = |\mathbb{E}_{\mathcal{D}}[P_n(A)] - \mathbb{E}_{\mathcal{U}}[P_n(A)]| \leq O(\ell^{-0.2k}) \leq \delta.\end{aligned}$$

Proof of Claim 18. We argue for $\mathcal{D}$, the same argument holds for $\mathcal{U}$. Write

$$|P_n(A) - P_k(A)| \leq \prod_{i=1}^{n} p_i \cdot \left| \sum_{i=k+1}^{n} S_i(A_1, \ldots, A_n) \right|.$$

If $A_1, \ldots, A_n$ are $(O(k), 0)$-wise independent, then, by Lemma 9,

$$\mathbb{E}[S_k(A_1, \ldots, A_n)^2] = \frac{(\sum_i \mathsf{Var}[A_i])^k}{k!} \leq \frac{\ell^{-0.6k}}{k!}.$$

Hence, under $(O(k), \varepsilon)$-wise independence,

$$\mathbb{E}[S_k(A_1, \ldots, A_n)^2] \leq \frac{\ell^{-0.6k}}{k!} + \varepsilon \mathrm{L}_1(S_k) \leq \frac{\ell^{-0.5k}}{k!}. \tag{36}$$

We now repeat the proof of Lemma 11 with $\sigma^2 = \ell^{-0.5}$ and $t = \ell^{0.2}$. The event G defined as

$$G = \left\{ |S_k(A)| \leq \frac{\ell^{-0.05k}}{\sqrt{k!}} \text{ and } |S_{k+1}(A)| \leq \frac{\ell^{-0.05(k+1)}}{\sqrt{(k+1)!}} \right\}$$

occurs with probability at least $1 - 2\ell^{-0.4k}$. As in the proof of Theorem 4, conditioned on G,

$$|P_n(A) - P_k(A)| \leq \sum_{i=k}^{n} |S_i(A)| \leq 2(6e\ell^{-0.05})^k. \tag{37}$$

Since $\mathbb{E}[A_i] = 0$ for all i and $\mathrm{L}_1(S_k) \leq \ell^{O(k)}$, by Equation (36), it follows that under $(O(k), \varepsilon)$-wise independence,

$$\mathbb{E}[P_k(A_1, \ldots, A_n)^2] \leq \sum_{i=0}^{k} \mathbb{E}[S_i(A_1, \ldots, A_n)^2] + \varepsilon \ell^{O(k)} = O(1). \tag{38}$$

Denote by $\neg G$ the complement of G. Write

$$\mathbb{E}[P_n(A) - P_k(A)] = \mathbb{E}[(P_n(A) - P_k(A))\mathbb{1}(G)] + \mathbb{E}[(P_n(A) - P_k(A))\mathbb{1}(\neg G)].$$

Equation (37) implies

$$|\mathbb{E}[(P_n(A) - P_k(A))\mathbb{1}(G)]| \leq 2(20\ell^{-0.25})^k.$$

It remains to bound the second term. Bound

$$|\mathbb{E}[(P_n(A) - P_k(A))\mathbb{1}(\neg G)]| \leq |\mathbb{E}[P_n(A)\mathbb{1}(\neg G)]| + |\mathbb{E}[P_k(A)\mathbb{1}(\neg G)]|$$

Note that $0 \leq P_n(A) \leq 1$ since it is the probability of an event. Also note that $\mathbb{E}[P_k(A)^2] = O(1)$ by Equation (38). So we can bound the RHS using Holder's inequalities by

$$|\mathbb{E}[\mathbb{1}(\neg G)]| + |\mathbb{E}[P_k(A)^2]^{1/2} \cdot \mathbb{E}[\mathbb{1}(\neg G)]^{1/2}| \leq O((\mathbb{E}[\mathbb{1}(\neg G)])^{1/2}) = O(\ell^{-0.2k}).$$

□

□

We are ready to prove the main theorem of this section.

Proof. The proof uses an hybrid argument. The $\mathcal{G}_{\mathcal{MR}}$ generator chooses $g_0 : [n] \to [m_0]$, and then $g'_1, \ldots, g'_T$ where $g'_t = [m_{t-1}] \times [n] \to [m_t]$ has error $\delta' = \delta/T$ and defines

$$g_t(i) = g'_t(g_{t-1}(i), i).$$

Let $h_0, h'_1, \ldots, h'_t$ be truly random hash functions with similar domains and ranges. For $0 \leq t, l \leq T$, define the hybrid family $\mathcal{G}^l_t = \{f^l_t : [m] \to [n]\}$ as follows: for $t = 0$ and every l,

$$f^l_0 = h_0,$$

and for $t > 0$ and every l,

$$f^l_t(i) = \begin{cases} g'_t(f^l_{t-1}(i), i) & \text{for } l < t, \\ h'_t(f^l_{t-1}(i), i) & \text{for } t \leq l. \end{cases}$$

For every l, let $\mathcal{G}^l = \mathcal{G}^l_T$. Thus, $\mathcal{G}^0 = \mathcal{G}_{\mathcal{MR}}$ and $\mathcal{G}^T = \mathcal{U}$. We will show by induction on $l \geq 1$ that

$$\left| \Pr_{f^l \in \mathcal{G}^l}[\forall\ i \in [n]\ f^l(i) \in S_i] - \Pr_{f^{l-1} \in \mathcal{G}^{l-1}}[\forall\ i \in [n]\ f^{l-1}(i) \in S_i] \right| \leq \delta'.$$

The desired bound then follows by the triangle inequality.

In the base case when $l = 1$, couple $\mathcal{G}^0$ and $\mathcal{G}^1$ by picking the same $g'_1, \ldots, g'_T$, and use them to define the function $f' : [m_1] \times [n] \to [m]$ so that

$$f^0(i) = f'(g_0(i), i),\ f^1(i) = f'(h_0(i), i).$$

For $i \in [n]$, define

$$S'_i = \{a \in [m_1] : f'(a, i) \in S_i\}.$$

Thus,

$$\begin{aligned} &\left| \Pr_{f^1 \in \mathcal{G}^1}[\forall\ i \in [n]\ f^1(i) \in S_i] - \Pr_{f^0 \in \mathcal{G}^0}[\forall\ i \in [n]\ f^0(i) \in S_i] \right| \\ &\qquad = \left| \Pr_{h_0}[\forall\ i \in [n]\ h_0(i) \in S'_i] - \Pr_{g_0}[\forall\ i \in [n]\ g_0(i) \in S'_i] \right| \leq \delta', \end{aligned}$$

by applying Lemma 16 with $k = O(\log(1/\delta'))$ and $\varepsilon = \delta' \cdot m_0^{-O(k)}$.

For the inductive case $l > 1$, couple $\mathcal{G}^l$ and $\mathcal{G}^{l-1}$ by picking the same $g'_{l+1}, \ldots, g'_T$, and pick $x \in [m_{l-1}]^n$ uniformly at random. There is a function $f' : [m_l] \times [n] \to [m]$ so that

$$f^l(i) = f'(h'_l(x_i, i), i),\ f^{l-1}(i) = f'(g'_l(x_i, i), i).$$

As before, define

$$S_i = \{a \in [m_l] : f'(a, i) \in S_i\}.$$

Hence,

$$\left| \Pr_{f^l \in \mathcal{G}^l}[\forall\ i \in [n]\ f^l(i) \in S_i] - \Pr_{f^{l-1} \in \mathcal{G}^{l-1}}[\forall\ i \in [n]\ f^{l-1}(i) \in S_i] \right|$$
$$= \left| \Pr_{h_l,x}[\forall\ i \in [n]\ h'_l(x_i, i) \in S'_i] - \Pr_{g_l,x}[\forall\ i \in [n]\ g'_l(x_i, i) \in S'_i] \right| \leq \delta,$$

by Lemma 17 with

$$k_{l-1} > 1,\ m_{l-1} \geq \log(1/\delta')^C,\ m_{l-1} \geq k_{l-1},\ m_{l-1}^{-k} \leq {\delta'}^C,\ \varepsilon_{l-1} \leq (m_l m_{l-1})^{-Ck}.$$

□

Acknowledgements

We thank Nati Linial, Raghu Meka, Yuval Peres, Dan Spielman, Avi Wigderson and David Zuckerman for helpful discussions. We thank an anonymous referee for pointing out an error in the statement of Theorem 4 in a previous version of the paper.

References

[ABI86] Noga Alon, Laszlo Babai, and Alon Itai. A fast and simple randomized parallel algorithm for the maximal independent set problem. *J. Algorithms 7(4)*, 1986.

[ASWZ96] Roy Armoni, Michael E. Saks, Avi Wigderson, and Shiyu Zhou. Discrepancy sets and pseudorandom generators for combinatorial rectangles. In *37th Annual Symposium on Foundations of Computer Science, FOCS '96*, pages 412–421, 1996.

[BCFM00] Andrei Z. Broder, Moses Charikar, Alan M. Frieze, and Michael Mitzenmacher. Min-wise independent permutations. *J. Comput. Syst. Sci.*, 60(3):630–659, 2000.

[BCM98] Andrei Z. Broder, Moses Charikar, and Michael Mitzenmacher. A derandomization using min-wise independent permutations. In *Randomization and Approximation Techniques in Computer Science, Second International Workshop, RANDOM'98*, pages 15–24, 1998.

[CG89] Benny Chor and Oded Goldreich. On the power of two-point based sampling. *J. Complexity 5(1)*, 1989.

[EGL+98] Guy Even, Oded Goldreich, Michael Luby, Noam Nisan, and Boban Velickovic. Efficient approximation of product distributions. *Random Struct. Algorithms*, 13(1):1–16, 1998.

[GKM15] Parikshit Gopalan, Daniel Kane, and Raghu Meka. Pseudorandomness via the discrete Fourier transform. In *Accepted to IEEE FOCS 2015*, 2015.

[GMR+12] Parikshit Gopalan, Raghu Meka, Omer Reingold, Luca Trevisan, and Salil P. Vadhan. Better pseudorandom generators from milder pseudorandom restrictions. In *53rd Annual IEEE Symposium on Foundations of Computer Science, FOCS'2012*, pages 120–129, 2012.

[Ind99] Piotr Indyk. A small approximately min-wise independent family of hash functions. In *Proceedings of the Tenth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 454–456, 1999.

[LLSZ97] Nathan Linial, Michael Luby, Michael E. Saks, and David Zuckerman. Efficient construction of a small hitting set for combinatorial rectangles in high dimension. *Combinatorica*, 17(2):215–234, 1997.

[Lu02] Chi-Jen Lu. Improved pseudorandom generators for combinatorial rectangles. *Combinatorica*, 22(3):417–434, 2002.

[Mul96] Ketan Mulmuley. Randomized geometric algorithms and pseudorandom generators. *Algorithmica*, 16(4/5):450–463, 1996.

[NN93] Joseph Naor and Moni Naor. Small-bias probability spaces: Efficient constructions and applications. *SIAM J. on Comput.*, 22(4):838–856, 1993.

[PS76] G Polya and G. Szego. *Problems and Theorems in Anaysis II.* Springer Classics in Mathematics, 1976.

[SSZZ99] Michael E. Saks, Aravind Srinivasan, Shiyu Zhou, and David Zuckerman. Low discrepancy sets yield approximate min-wise independent permutation families. In *Third International Workshop on Randomization and Approximation Techniques in Computer Science RANDOM'99*, pages 11–15, 1999.

[Ste04] J. Michael Steele. *The Cauchy-Schwarz Master Class.* Cambridge University Press, 2004.

A Missing Proofs

We give the proof of Fact B which states that over the reals, if $S_k(a) = S_{k+1}(a) = 0$ for $k > 0$ then $S_\ell(a) = 0$ for all $\ell \geq k$.

For a univariate polynomial $p(\xi)$ and a root $y \in \mathbb{R}$ of p, denote by $\mathsf{mult}(p, y)$ the multiplicity of the root y in p. We use the following property of polynomials $p(\xi)$ with real roots [PS76], which can be proved using the interlacing of the zeroes of $p(\xi)$ and $p'(\xi)$: If $\mathsf{mult}(p', y) \geq 2$ then $\mathsf{mult}(p, y) \geq \mathsf{mult}(p', y) + 1$.

Proof of Fact B. Let

$$p(\xi) = \prod_{i \in [n]} (\xi + b_i) = \sum_{k=0}^{n} \xi^k S_{n-k}(b).$$

Consider $p^{(n-k-1)}(\xi)$ which is the $(n-k-1)^{th}$ derivative of $p(\xi)$. Since $S_k(b) = S_{k+1}(b) = 0$ for $k > 0$, it follows that ξ^2 divides $p^{(n-k-1)}(\xi)$ and hence $\mathsf{mult}(p^{(n-k-1)}, 0) \geq 2$. Applying the above fact $n-k-1$ times, we get $\mathsf{mult}(p, 0) \geq n-k+1$ so $S_n(b) = \ldots = S_k(b) = 0$. □

The next Theorem is a routine extension of the result of [SSZZ99] to large ℓ.

Theorem 19. *[SSZZ99] Let $\mathcal{G} : \{0,1\}^r \to [m]^n$ be a* PRG *for combinatorial rectangles with error ε. The resulting family $\{g_y : y \in \{0,1\}^r\}$ of hash functions is approximately ℓ-minima-wise independent with error at most $\varepsilon\binom{m}{\ell}$.*

Proof of Theorem 19. Fix $S \subseteq [n]$ and a sequence $T = (t_1, \ldots, t_\ell)$ of ℓ distinct elements from S. The event

$$g(t_1) < \cdots < g(t_\ell) < \min g(S \setminus T)$$

can be viewed as the disjoint union of $\binom{m}{\ell}$ events by fixing the set $A = \{a_1 < \ldots < a_\ell\}$ that T maps to. The indicator $\mathbb{1}_A$ of the event

$$g(t_1) = a_1, \ldots,\ g(t_\ell) = a_\ell,\ g(S \setminus T) > a_\ell$$

is a combinatorial rectangle: Define

$$\begin{aligned} f_i(x_i) &= 1 \text{ for } i \notin S \\ f_i(x_i) &= \mathbb{1}(x_i = a_j) \text{ for } i = t_j \in T \\ f_i(x_i) &= \mathbb{1}(x_i > a_\ell) \text{ for } i \in S \setminus T \end{aligned}$$

and

$$f_A(x_1, \ldots, x_n) = \bigwedge_{i \in [n]} f_i(x_i).$$

Since $g(i) = x_i$, it follows that $\mathbb{1}_A(g) = f_A(x)$. Further, choosing $h \in \mathcal{U}$ is equivalent to choosing $x \in [m]^n$ uniformly at random. Hence,

$$\begin{aligned} &\Pr_{g \in \mathcal{G}}[g(t_1) < \cdots < g(t_\ell) < \min g(S \setminus T)] \\ &\qquad = \sum_A \mathbb{E}_{y \in \{0,1\}^r}[f_A(\mathcal{G}(y))] \\ &\qquad = \sum_A (\mathbb{E}_{h \in \mathcal{U}}[\mathbb{1}_A(h)] \pm \varepsilon) \\ &\qquad = \Pr_{h \in \mathcal{U}}[h(t_1) < \cdots < h(t_\ell) < \min h(S \setminus T)] \pm \binom{m}{\ell}\varepsilon. \end{aligned}$$

□

Finally we discuss how to generate the (k, ε)-wise independent distributions on $[m]^n$ with seed length $O(\log\log(n) + k\log(m) + \log(1/\varepsilon))$. We claim that it suffices to take a $k' = k\log(m)$-wise ε-independent string of length $n' = n\log(m)$. Naor and Naor [NN93] showed that such distributions can be generated using seed-length $O(\log\log(n) + k\log(m) + \log(1/\varepsilon))$. We can also assume that every co-ordinate is uniformly random in $[m]$ by adding the string $(a, \ldots, a)$ where $a \in [m]$ is chosen randomly.